



Insect community responses to prescribed fire & White-nose Syndrome in eastern deciduous forests


Luke E. Dodd¹, Shelby A. Fulton¹, Rachael E. Griffiths¹, & Lynne K. Rieske²

¹Department of Biological Sciences, Eastern Kentucky University, Richmond, KY (luke.dodd@eku.edu)

²Department of Entomology, University of Kentucky, Lexington, KY


Introduction

- Insectivorous bats are important predators; studies suggest bats can exert top-down pressure on prey communities^{1,2}
- Ecosystem services provided by bats in North America total in the billions of dollars³
- White-nose Syndrome (WNS) is an emergent disease responsible for the deaths of millions of bats in eastern North America
- WNS has destabilized bat community structure and relaxed niche partitioning⁴




Objectives & Study Design



- We are elucidating the multi-trophic impacts of WNS in a fire-managed landscape
- Mammoth Cave National Park; WNS first detected in Jan 2013



Methods & Results




Acoustic ID of bats



Light-trapping surveys




Estimates of insect abundance



Expected Outcomes & Implications

- Due to relaxed niche partitioning post-WNS, bats will adopt non-traditional foraging niches and thus be exposed to a new composition of available prey. The dramatic mortality associated with WNS may lead to reduced competition for prey resources among sympatric bat species.



- Diets may expand to include additional insect orders or narrow to rely more on preferred insect orders

Thanks!

Especially Tracy Culbertson,
Clint Rose, and Jennifer Winters!

Collaborators

Matt Dickinson, Mike Lacki, Nick Skowronski, Steve Thomas, Rick Toomey

Funding

JFSP 10-1-06-1, JFSP 14-1-05-22

Lit Cited

1. Kalka et al. 2008. Science 320: 71.
2. Williams-Guillen et al. 2008. Science 320: 70.
3. Boyles et al. 2011. Science 332: 41-42.
4. Jachowski et al. 2014. Diversity and Distributions 20: 1002–1015.